arXiv:math/0405131v1 [math.RA] 7 May 2004

Algebras of non-Archimedean measures on groups

S. V. Lüdkovsky

7 May 2004

Quasi-invariant measures with values in non-Archimedean fields on a group of diffeomorphisms were constructed for non-Archimedean manifolds M in [Lud96, Lud99t]. On non-Archimedean loop groups and semigroups they were provided in [Lud98s, Lud00a, Lud02b]. A Banach space over a local field also serves as the additive group and quasi-invariant measures on it were studied in [Lud03s2, Lud96c].

This article is devoted to the investigation of properties of quasi-invariant measures with values in non-Archimedean fields that are important for analysis on topological groups and for construction of irreducible representations. The following properties are investigated:

(1) convolutions of measures and functions,

(2) continuity of functions of measures,

(3) non-associative algebras generated with the help of quasi-invariant measures. The theorems given below show that many differences appear to be between locally compact and non-locally compact groups. The groups considered below are supposed to have structure of Banach manifolds over the corresponding fields.

1. Definitions. (a). Let G be a Hausdorff separable topological group. A tight measure μ on $Af(G,\mu)$ with values in a non-Archimedean field $\mathbf{F}$ is called left-quasi-invariant (or right) relative to a dense subgroup H of G, if $\mu_\phi(*)$ (or $\mu^\phi(*)$) is equivalent to $\mu(*)$ for each $\phi \in H$, where $Bco(G)$ is the algebra of all clopen subsets of G, $Af(G,\mu)$ denotes its completion by μ, $\mu_\phi(A) := \mu(\phi^{-1}A)$, $\mu^\phi(A) := \mu(A\phi^{-1})$ for each $A \in Af(G,\mu)$, $\rho_\mu(\phi,g) := \mu_\phi(dg)/\mu(dg) \in L(G, Af(G,\mu), \mu, \mathbf{F})$ (or $\tilde{\rho}_\mu(\phi,g) := \mu^\phi(dg)/\mu(dg)$) denotes a left (or right) quasi-invariance factor, $\mathbf{F}$ is a non-Archimedean field complete relative to its uniformity and such that $\mathbf{K}_s \subset \mathbf{F}$. We assume that a uniformity τ_G on G is such that $\tau_G|H \subset \tau_H$, (G,τ_G) and (H,τ_H) are complete. We suppose also that there exists an open base in $e \in H$ such that their closures in G are compact (such pairs exist for loop groups and groups of diffeomorphisms and Banach-Lie groups). We denote by $M_l(G,H)$ (or $M_r(G,H)$) a set of left-(or right) quasi-invariant tight measures on G relative to H with a finite norm $\|\mu\| < \infty$.

(b). Let $L_H(G,\mu,\mathbf{F})$ denotes the Banach space of functions $f : G \to \mathbf{F}$ such that $f_h(g) \in L(G,\mu,\mathbf{F})$ for each $h \in H$ and

$$\|f\|_{L_H(G,\mu,\mathbf{F})} := \sup_{h\in H} \|f_h\|_{L(G,\mu,\mathbf{F})} < \infty,$$

where $\mathbf{F}$ is a non-Archimedean field for which $\mathbf{K}_s \subset \mathbf{F}$, $f_h(g) := f(h^{-1}g)$ for each $g \in G$. For $\mu \in M_l(G,H)$ and $\nu \in M(H)$ let

$$(\nu * \mu)(A) := \int_H \mu_h(A)\nu(dh) \text{ and } (q\tilde{*}f)(g) := \int_H f(hg)q(h)\nu(dh)$$

be convolutions of measures and functions, where $M(H)$ is the space of tight measures on H with a finite norm, $\nu \in M(H)$ and $q \in L(H,\nu,\mathbf{F})$.

2. Lemma. *The convolutions are continuous* $\mathbf{F}$*-linear mappings*

$$* : M(H) \times M_l(G,H) \to M_l(G,H) \textit{ and}$$

$$\tilde{*} : L(H,\nu,\mathbf{F}) \times L_H(G,\mu,\mathbf{F}) \to L_H(G,\mu,\mathbf{F}).$$

Proof. This follows from Theorem 7.16 [Roo78] and estimates

$$\|\nu * \mu\| \le \|\nu\| \times \|\mu\|, \ \|q \tilde{*} f\|_{L_H(G,\mu,\mathbf{F})} \le \|q\|_{L(H,\nu,\mathbf{F})} \times \|f\|_{L_H(G,\mu,\mathbf{F})},$$

since $\rho_\mu(h,g) \in L(H \times G, \nu \times \mu, \mathbf{F})$.

3. Lemma. *For $\mu \in M_l(G,H)$ the translation map $(q,f) \to f_q(g)$ is continuous from $H \times L_H(G,\mu,\mathbf{F})$ into $L_H(G,\mu,\mathbf{F})$.*

Proof. In view of Lemma 7.10 and Theorem 7.12 [Roo78] for each $\epsilon > 0$ the set $\{x : |f(x)|N_\mu(x) \ge \epsilon\}$ is $Af(G,\mu)$-compact and f is $Af(G,\mu)$-continuous. The embedding of H into G is compact (see §1), hence for each $q \in H$ there exists V clopen in H and such that $q^{-1}V$ is a subgroup of H with $cl_G q^{-1}V$ compact in G, where $cl_G(A)$ denotes the closure of a subset A in G. The product of compact subsets in G is compact in G, hence $f_q(g) \in L_H(G,\mu,\mathbf{F})$ and $(q,f) \mapsto f_q(g)$ is the continuous mapping, since the restriction of the $Bco(G)$ and the $Af(G,\mu)$-topologies onto X_ϵ coincide, $\|f_q\|_{L_H(\mu)} = \|f\|_{L_H(\mu)}$ for each $q \in H$ (see §1.(b)).

4. Proposition. *For a probability measure $\mu \in M(G)$ there exists an approximate unit which is a sequence of nonzero continuous functions $\psi_i : G \to \mathbf{F}$ such that $\int_G \psi_i(g)\mu(dg) = 1$ and for each neighbourhood $U \ni e$ in G there exists i_0 such that $supp(\psi_i) \subset U$ for each $i > i_0$.*

Proof. A group G has a countable base of neighbourhoods of $e \in G$. A measure μ is quasi-invariant, hence $\|U\|_\mu > 0$ for each neighbourhood $U \ni e$, μ is the tight measure, hence there exists a system of neighbourhoods $\{U_i : U_i \ni e \forall i\}$, $\bigcap_i U_i = \{e\}$, $U_i \supset U_{i+1}$ for each i, $supp(\psi_i) \subset U_i$. Choose ψ_i such that $\int_G \psi_i(g)\mu(dg) = 1$ for each i.

5. Proposition. *If $(\psi_i : i \in \mathbf{N})$ is an approximate unit in H relative to a probability measure $\nu \in M(H)$, then $\lim_{i\to\infty} \psi_i * f = f$ in the $L_H(G,\mu,\mathbf{F})$ norm, where $\mu \in M_l(G,H)$, $f \in L_H(G,\mu,\mathbf{F})$.*

Proof. In view of Theorem 7.12 [Roo78] for each $\epsilon > 0$ and each i there exists a finite number of $h_j \in H$, $j = 1,...,n$, $n \in \mathbf{N}$, such that

$\bigcup_{j=1}^{n} h_jU_i \supset X_{\epsilon,f} \times V$, where $\{x : |f(x)|N_\mu(x) \geq \epsilon\} =: X_{\epsilon,f}$, V is a clopen neighbourhood of $\xi \in H$ which can be chosen such that $\xi^{-1}V$ is a subgroup of H, $cl_G(\xi^{-1}V)$ is compact in G,

$(\psi_i * f)(g) = \int_{h \in U_i} \psi_i(h) f(hg) \nu(dh)$ and
$\sup_{g\in G} |f_\xi(g) - (\psi_i * f_\xi)(g)| N_\mu(g) \leq \sup_{g \in G}[\sup_{h \in X_{\epsilon,f_\xi}} |\psi_i(h)||f_\xi(hg) - f_\xi(g)|N_\mu(g) + \epsilon\|\psi_i\|_\nu \|G\|_\mu]$,
hence $\lim_{i\to\infty}(\psi_i * f) = f$ in $L_H(G,\mu,\mathbf{F})$-norm.

6. Lemma. *Suppose* $g \in L_H(G,\mu,\mathbf{F})$ *and* $(g^x|_H) \in L(H,\nu,\mathbf{F})$ *for each* $x \in G$, $f \in L(H,\nu,\mathbf{F})$, *where* $g^x(y) := g(yx)$ *for each* x *and* $y \in G$. *Let* μ *and* ν *be probability measures,* $\mu \in M_l(G,H)$, $\nu \in M(H)$. *Then* $f\tilde{*}g \in L_H(G,\mu,\mathbf{F})$ *and there exists a function* $h : G \to \mathbf{F}$ *such that* $h|_H$ *is continuous,* $h = f\tilde{*}g$ μ*-a.e. on* G *and* h *vanishes at* ∞ *on* G.

Proof. In view of Fubini theorem we have

$$\|f\tilde{*}g\|_{L_H(G,\mu,\mathbf{F})} = \sup_{h\in H, z \in G} |\int_H f(y)g(yz)\nu(dy)|N_\mu(z) \leq$$

$$\|g(z)\|_{L_H(G,\mu,\mathbf{F})}\|f\|_{L(H,\nu,\mathbf{F})}$$

The equation $\alpha_f(\phi) := \int_H f(y)\phi(y)\nu(dy)$ defines a continuous linear functional on $L^\infty(H,\nu,\mathbf{F}) := \{\phi : H \to \mathbf{F} : \phi$ is $(Af(H,\nu), Bco(\mathbf{F}))$-measurable, $\|\phi\|_\infty := ess_\nu - \sup_{x\in H}|\phi(x)| < \infty\}$. In view of Lemma 3 the function $\alpha_f(g^{(qx)^{-1}}) =: v(qx) =: w(q,x)$ of two variables q and x is continuous on $H \times H$ for $q, x \in H$, since the mapping $(q,x) \mapsto (qx)^{-1}$ is continuous from $H \times H$ into H. By Theorem 7.16 [Roo78]

$$\int_G v(y)\psi(y)\mu(dy) = \int_G \int_H f(y)g(yx)\psi(x)\nu(dy)\mu(dx) =$$

$$\int_H f(y)[\int_G g(yx)\psi(x)\mu(dx)]\nu(dy)$$

for each $\psi \in L^\infty(G,\mu,\mathbf{F})$. From this it follows, that $\mu(\{y : h(y) \neq (f\tilde{*}g)(y), y \in G\}) = 0$, since h and $(f\tilde{*}g)$ are μ-measurable functions due to Fubini theorem and the continuity of the composition and the inversion in a topological group. In view of Theorem 7.12 [Roo78] for each $\epsilon > 0$ there are compact subsets $C \subset H$ and $D \subset G$ and functions $f' \in L(H, \nu, \mathbf{F})$ and $g' \in L_H(G, \mu, \mathbf{F})$ with closed supports $supp(f') \subset C$, $supp(g') \subset D$ such that $cl_G CD$ is compact in G,

$$\|f' - f\|_{L(H,\nu,\mathbf{F})} < \epsilon \text{ and } \|g' - g\|_{L_H(G,\mu,\mathbf{F})} < \epsilon,$$

since by the supposition of §1 the group H has the base B_H of its topology τ_H, such that the closures $cl_G V$ are compact in G for each $V \in \mathsf{B}_H$. From the inequality

$$|h'(x) - h(x)| \le (\|f\|_{L(H,\nu,\mathbf{F})} + \epsilon)\epsilon + \epsilon \|g\|_{L_H(G,\mu,\mathbf{F})}$$

it follows that for each $\delta > 0$ there exists a compact subset $K \subset G$ with $|h(x)| < \delta$ for each $x \in G \setminus K$, where $h'(x^{-1}) := \alpha_{f'}(g'^x)$.

7. Proposition. *Let* $A, B \in Af(G, \mu)$, μ *and* ν *be probability measures,* $\mu \in M_l(G, H)$, $\nu \in M(H)$. *Then the function* $\zeta(x) := \mu(A \cap xB)$ *is continuous on* H *and* $\nu(yB^{-1} \cap H) \in L(H, \nu, \mathbf{F})$. *Moreover, if* $\|A\|_\mu \|B\|_\mu > 0$, $\mu(\{y \in G : yB^{-1} \cap H \in Af(H, \nu)$ *and* $\|yB^{-1} \cap H\|_\nu > 0\}) > 0$, *then* $\zeta(x) \ne 0$ *on* H.

Proof. Let $g_x(y) := Ch_A(y) Ch_B(x^{-1}y)$, then $g_x(y) \in L_H(G, \mu, \mathbf{F})$, where $Ch_A(y)$ is the characteristic function of A. In view of Propositions 4 and 5 there exists $\lim_{i\to\infty} \psi_i * g_x = g_x$ in $L_H(G, \mu, \mathbf{F})$. In view of Lemma 6 $\zeta(x)|_H$ is continuous. There is the following inequality:

$$1 \ge |\int_H \mu(A \cap xB)\nu(dx)| = |\int_H \int_G Ch_A(y) Ch_B(x^{-1}y)\mu(dy)\nu(dx)|.$$

In view of Theorem 7.16 [Roo78] there exists

$$\int_H Ch_B(x^{-1}y)\nu(dy) = \nu((yB^{-1}) \cap H) \in L(G, \mu, \mathbf{F}), \text{ hence}$$

$$\int_H \mu(A \cap xB)\nu(dx) = \int_G \nu(yB^{-1} \cap H) Ch_A(y)\mu(dy).$$

8. Corollary. *Let* $A, B \in Af(G,\mu)$, $\nu \in M(H)$ *and* $\mu \in M_l(G,H)$ *be probability measures. Then denoting* $Int_H V$ *the interior of a subset* V *of* H *with respect to* τ_H, *one has*

(i) $Int_H(AB) \cap H \ne \emptyset$, *when*

$$\|\{y \in G : \|yB \cap H\|_\nu > 0\}\|_\mu > 0;$$

(ii) $Int_H(AA^{-1}) \ni e$, *when*

$$\|\{y \in G : \|yA^{-1} \cap H\|_\nu > 0\}\|_\mu > 0.$$

Proof. $AB \cap H \supset \{x \in H : \|(A \cap xB^{-1})\|_\mu > 0\}$.

9. Corollary. *Let* $G = H$. *If* $\mu \in M_l(G,H)$ *is a probability measure, then* G *is a locally compact topological group.*

Proof. Let us take $\nu = \mu$ and $A = C \cup C^{-1}$, where C is a compact subset of G with $\|(C)\|_\mu > 0$, whence $\|(yA)\|_\mu > 0$ for each $y \in G$ and inevitably $Int_G(AA^{-1}) \ni e$.

10. Corollary. *Let* $\mu \in M_l(G,H)$, $\rho_\mu(h,z) \in L(H,\nu,\mathbf{F}) \times L(G,\mu,\mathbf{F})$, *then* $\rho_\mu(h,z)$ *is continuous* $\nu \times \mu$*-a.e. on* $H \times G$.

Proof. In view of the cocycle condition
$\rho_\mu(\phi\psi, g) := \mu_{\phi\psi}(dg)/\mu(dg) = (\mu_{\phi\psi}(dg)/\mu_\phi(dg))(\mu_\phi(dg)/\mu(dg))$
$= \rho_\mu(\psi, \phi^{-1}g)\rho_\mu(\phi, g)$
on ρ_μ and Corollary 8 above, Theorem 7.12 [Roo78] for each $\epsilon > 0$ the quasi-invariance factor $\rho_\mu(h,g)$ is continuous on $H \times G_\epsilon$, but G_ϵ is neighbourhood of e in G, where $G_\epsilon := \{g \in G : N_\mu(g) \ge \epsilon\}$. Since H is dense in G and G is separable, then $\bigcup_{j=1}^\infty h_j G_\epsilon = G$, where $\{h_j : j \in \mathbf{N}\}$ is a countable subset in G. Therefore, $\rho_\mu(h,z)$ is continuous $\nu \times \mu$-a.e. on $H \times G$.

11. Corollary. *Let* G *be a locally compact group,* $\rho_\mu(h,z) \in L(G \times G, \mu \times \mu, \mathbf{F})$, *then* $\rho_\mu(h,z)$ *is* $\mu \times \mu$*-a.e. continuous on* $G \times G$.

Proof. It follows from Corollaries 9 and 10.

12. Remark. The latter two corollaries show, that the condition of continuity of $\rho_\mu(h, z)$ imposed in [Lud03s2, Lud96c] is not restrictive.

13. Lemma. *Let* $\mu \in M_l(G, H)$ *be a probability measure and* G *be non-locally compact. Then* $\|H\|_\mu = 0$.

Proof. This follows from Theorem 3.13 [Lud96c] above and the proof of Lemma 2, since the embedding $T_eH \hookrightarrow T_eG$ is a compact operator in the non-Archimedean case and a tight measure μ on G induces a tight measure on a neighbourhood V of 0 in T_eG such that V is topologically homeomorphic to a clopen subgroup U in G (see also papers about construction of quasi-invariant measures on the considered here groups [Lud96, Lud99t, Lud98s, Lud00a, Lud02b]).

14. Let (G, τ_G) and (H, τ_H) be a pair of topological non-locally compact groups G, H (Banach-Lie, Frechet-Lie or groups of diffeomorphisms or loop groups) with uniformities τ_G, τ_H such that H is dense in (G, τ_G) and there is a probability measure $\mu \in M_l(G, H)$ with continuous $\rho_\mu(z, g)$ on $H \times G$. Also let X be a Banach space over $\mathbf{F}$ and $IS(X)$ be the group of isometric $\mathbf{F}$-linear automorphisms of X in the topology inherited from the Banach space $L(X)$ of all bounded $\mathbf{F}$ linear operators from X into X.

Theorem. (1). *If* $T : G \to IS(X)$ *is a weakly continuous representation, then there exists* $T' : G \to IS(X)$ *equal* μ*-a.e. to* T *and* $T'|_{(H,\tau_H)}$ *is strongly continuous.*

(2). *If* $T : G \to IS(X)$ *is a weakly measurable representation and* X *is of separable type* $c_0(\mathbf{F})$ *over* $\mathbf{F}$*, then there exists* $T' : G \to IS(X)$ *equal to* T μ*-a.e. and* $T'|_{(H,\tau_H)}$ *is strongly continuous.*

Proof. Take $\mathcal{K}(G) := (I) \cup L(G, \mu, \mathbf{F})$, where I is the unit operator on $L(G, \mu, \mathbf{F})$. Then we can define

$$A_{(\lambda e + a)_h} := \lambda I + \int_G a_h(g)[\rho_\mu(h, g)]T_g \mu(dg),$$

where $a_h(g) := a(h^{-1}g)$. Then

$$|(A_{(\lambda e+a)_h} - A_{\lambda e+a}\xi, \eta)| \le \sup_{g\in G}[|a_h(g)\rho_\mu(h,g) - a(g)|\ |T_g\xi| N_\mu(g)],$$

hence A_{a_h} is strongly continuous with respect to $h \in H$, that is,

$$\lim_{h\to e} |A_{a_h}\xi - A_a\xi| = 0.$$

Denote $A_{a_h} = T'_h A_a$, so $T'_h\xi = A_{a_h}\xi_0$, where $\xi = A_a\xi_0$, $a \in L(G,\mu,\mathbf{F})$. Whence

$$\begin{aligned} T'{}_h\xi = A_{a_h}\xi_0 &= \int_G a_h(g)T_g\xi_0\rho_\mu(h,g)\mu(dg) \\ &= T_h\int_G a(z)T_z\xi_0\mu(dz) = T_h\xi, \end{aligned}$$

hence $|T'{}_h\xi| = |\xi|$ for each $h \in H$. Therefore, T'_h is uniquely extended to an isometric operator on the Banach space $X' \subset X$. In view of Lemma 10, $\|H\|_\mu = 0$. Hence T' may be considered equal to T μ-a.e. Then a space $sp_{\mathbf{F}}[A_{a_h} : h \in H]$ is evidently dense in X, since

$$A_{a_h}\xi = \int_G a_h(g)T_g\rho_\mu(h,g)\mu(dg)\xi =$$

$$T_h\int_G a(g)T_g\mu(dg)\xi.$$

For proving the second statement let
$\mathsf{R} := [\xi : A_a\xi = 0$ for each $a \in L(G,\mu,\mathbf{F})]$. If $\eta \in X^*$ and

$$\eta(A_a\xi) = \eta(\int_G a(g)T_g\xi\mu(dg)) = \eta(\int_G a(g)T'{}_g\xi\mu(dg))$$

for each $a(g) \in L(G,\mu,\mathbf{F})$, then $\eta(T_g\xi) = \eta(T'_g\xi)$ for μ-almost all $g \in G$, where X^* denotes the topological dual space of all $\mathbf{K}$-linear functionals $f : X \to \mathbf{K}$. Suppose that $\{\xi^*_n : n \in \mathbf{N}\}$ is an orthonormal system in X^* separating points of X. It exists, since by the supposition of this theorem $X = c_0(\mathbf{F})$. If $\xi \in X$, then

$$\xi^*_m(\int_G a(g)T_g\xi\mu(dg)) = 0$$

for each $g \in G \setminus S_m$, where $\|S_m\|_\mu = 0$. Therefore, $\xi^*_m(T_g\xi) = 0$ for each $m \in \mathbf{N}$, if $g \in G \setminus S$, where $S := \bigcup_{m=1}^\infty S_m$. Hence $T_g\xi = 0$ for each $g \in G \setminus S$, consequently, $\xi = 0$. Consider the embedding $X \hookrightarrow X^*$ with the help of the standard orthnormal basis $\{e_j : j\}$ in X over $\mathbf{F}$. Therefore, let $\{\xi_m : m \in \mathbf{N}\} \subset X \hookrightarrow X^*$. Then $\xi^*_m(T_g\xi_n) = \xi^*_m(T'_g\xi_n)$ for each $g \in G \setminus \gamma_{n,m}$, where $\|\gamma_{n,m}\|_\mu = 0$, ξ^*_m is the image of ξ_m under this standard embedding $X \hookrightarrow X^*$. Hence $\xi^*_m(T_g\xi_n) = \xi^*_m(T'_g\xi_n)$ for each $n, m \in \mathbf{N}$ and each $g \in G \setminus \gamma$, where $\gamma := \bigcup_{n,m} \gamma_{n,m}$ and $\|\gamma\|_\mu = 0$, since in $L(G, \mu, \mathbf{F})$ the family of all step functions is dense. Therefeore, $\mathsf{R} = 0$.

15. Definition and note. Let $\{G_i : i \in \mathbf{N_o}\}$ be a sequence of topological groups such that $G = G_0$, $G_{i+1} \subset G_i$ and G_{i+1} is dense in G_i for each $i \in \mathbf{N_o}$ and their topologies are denoted τ_i, $\tau_i|_{G_{i+1}} \subset \tau_{i+1}$ for each i, where $N_o := \{0, 1, 2, ...\}$. Suppose that these groups are supplied with $\mathbf{F}$-valued probability quasi-invariant measures μ^i on G_i relative to G_{i+1}. For example, such sequences exist for groups of diffeomorphisms or loop groups considered in previous papers [Lud96, Lud99t, Lud98s, Lud00a, Lud02b]). Let $L_{G_{i+1}}(G_i, \mu^i, \mathbf{F})$ denotes a Banach subspace of $L(G_i, \mu^i, \mathbf{F})$ as in §1(b). Let $\tilde{L}(G_{i+1}, \mu^{i+1}, L(G_i, \mu^i, \mathbf{F})) =: H_i$ denotes the completion of the subspace of $L(G_i, \mu^i, \mathbf{F})$ of all elements f such that

$$\|f\|_i := \max[\|f^2\|^{1/2}{}_{L(G_i,\mu^i,\mathbf{F})}, \|f\|'_i] < \infty, \text{ where}$$

$$\|f\|'_i := [\sup_{x \in G_i, y \in G_{i+1}} |f(y^{-1}x)|^2 N_{\mu^i}(x) \max(1, N_{\mu^{i+1}}(y))]^{1/2}.$$

Evidently H_i are Banach spaces over $\mathbf{F}$. Let

$$f^{i+1} * f^i(x) := \int_{G_{i+1}} f^{i+1}(y) f^i(y^{-1}x) \mu^{i+1}(dy)$$

denotes the convolution of $f^i \in H_i$.

16. Lemma. *The convolution* $* : H_{i+1} \times H_i \to H_i$ *is the continuous* $\mathbf{F}$*-bilinear mapping.*

Proof. From the definitions we have:
$\|(f^{i+1}*f^i)^2\|^{1/2}_{L(G_i,\mu^i,\mathbf{F})}=\sup_{x\in G_i}|(f^{i+1}*f^i)(x)|N^{1/2}_{\mu^i}(x)$
$\le \sup_{x\in G_i,y\in G_{i+1}}[|f^{i+1}(y)|N^{1/2}_{\mu^{i+1}}(y)][|f^i(y^{-1}x)|N^{1/2}_{\mu^i}(x)N^{1/2}_{\mu^{i+1}}(y)]$
$=\|(f^{i+1})^2\|^{1/2}_{L(G_{i+1},\mu^{i+1},\mathbf{F})}\|f^i\|'_i$ and
$\|f^{i+1}*f^i\|'_i=[\sup_{x\in G_i,y\in G_{i+1}}|(f^{i+1}*f^i)(y^{-1}x)|^2N_{\mu^i}(x)N_{\mu^{i+1}}(y)]^{1/2}$
$\le[\sup_{x\in G_i,y\in G_{i+1},z\in G_{i+1}}|f^{i+1}(z)|^2N_{\mu^{i+1}}(z)|f^i(y^{-1}z^{-1}x)|^2N_{\mu^i}(x)N_{\mu^{i+1}}(y)N_{\mu^{i+1}}(z)]^{1/2}$
$\le\|(f^{i+1})^2\|^{1/2}_{L(G_{i+1},\mu^{i+1},\mathbf{F})}\|f^i\|'_i$,
since from $A\ni y$ and $B\ni z$ for $A,B\in Af(G_{i+1},\mu^{i+1})$ it follows that $AB\ni yz$ and $AB\in Af(G_{i+1},\mu^{i+1})$, which follows from
$\mu^{i+1}(AB)=\int_{A\ni a}\mu^{i+1}(aB)\mu^{i+1}(da)=\int_{B\ni b}\int_{A\ni a}\mu^{i+1}(adb)\mu^{i+1}(da)$, so that $\|A\|_{\mu^{i+1}}\le\|G_{i+1}\|_{\mu^{i+1}}=1$, hence $N_{\mu^{i+1}}(z)\le 1$ for each $z\in G_{i+1}$ for the probability measure μ^{i+1}. Therefore, $\|f^{i+1}*f^i\|_i\le\|f^{i+1}\|_{i+1}\|f^i\|_i$.

17. Definition. Let $c_0(\{H_i: i\in\mathbf{N_o}\})=:H$ be the Banach space consisting of elements $f=(f^i: f^i\in H_i, i\in\mathbf{N_o})$, for which $\lim_{i\to\infty}\|f^i\|_i=0$, where

$$\|f\|:=\sup_{i=0}^{\infty}\|f^i\|_i<\infty.$$

For elements f and $g\in H$ their convolution is defined by the formula: $f\star g:=h$ with $h^i:=f^{i+1}*g^i$ for each $i\in\mathbf{N_o}$. Let $*: H\to H$ be an involution such that $f^*:=(f^{j\wedge}: j\in\mathbf{N_o})$, where $f^{j\wedge}(y_j):=f^j(y_j^{-1})$ for each $y_j\in G_j$, $f:=(f^j: j\in\mathbf{N_o})$.

18. Lemma. *H is a non-associative non-commutative Banach algebra with involution $*$, that is $*$ is $\mathbf{F}$-bilinear and $f^{**}=f$ for each $f\in H$.*

Proof. In view of Lemma 16 the convolution $h=f\star g$ in the Banach space H has the norm $\|h\|\le\|f\|\,\|g\|$, hence is a continuous mapping from $H\times H$ into H. From its definition it follows that the convolution is $\mathbf{F}$-bilinear. It is non-associative as follows from the computation of i-th terms of $(f\star g)\star q$ and $f\star(g\star q)$, which are $(f^{i+2}*g^{i+1})*q^i$ and $f^{i+1}*(g^{i+1}*q^i)$ respectively, where f, g and $q\in H$. It is non-commutative, since there are f and $g\in H$ for which $f^{i+1}*g^i$ are not equal to $g^{i+1}*f^i$. From $f^{j\wedge\wedge}(y_j)=f^j(y_j)$ it follows that $f^{**} = (f^*)^* = f$.

19. Note. In general $(f \star g^*)^* \ne g \star f^*$ for f and $g \in H$, since there exist f^j and g^j such that $g^{j+1} * (f^j)^* \ne (f^{j+1} * (g^j)^*)^*$. If $f \in H$ is such that $f^j|_{G_{j+1}} = f^{j+1}$, then

$$((f^{j+1})^* * f^j)(e) = \int_{G_{j+1}} (f^{j+1}(y))^2 \mu^{j+1}(dy), \text{ hence}$$

$$|((f^{j+1})^* * f^j)(e)| \le \|(f^{j+1})^2\|^{1/2}_{L(G_{j+1},\mu^{j+1},\mathbf{F})} \le \|f^{j+1}\|_{j+1},$$

where $j \in \mathbf{N_o}$.

20. Definition. Consider the standard Banach space $c_0(\mathbf{F})$ over the field $\mathbf{F}$ as a Banach algebra with the convolution $\alpha \star \beta = \gamma$ such that $\gamma^i := \alpha^{i+1}\beta^i$, where $\alpha := (\alpha^i : \alpha^i \in \mathbf{F}, i \in \mathbf{N_o})$, α, β and $\gamma \in c_0(\mathbf{F})$.

21. Note. The algebra $c_0(\mathbf{F})$ has two-sided ideals $J_i := \{\alpha \in c_0(\mathbf{F}) : \alpha^j = 0 \text{ for each } j > i\}$, where $i \in \mathbf{N_o}$. That is, $J \star c_0(\mathbf{F}) \subset J$ and $c_0(\mathbf{F}) \star J = J$ and J is the $\mathbf{F}$-linear subspace of $c_0(\mathbf{F})$, but $J \star c_0(\mathbf{F}) \ne J$. There are also right ideals, which are not left ideals: $K_i := \{\alpha \in c_0(\mathbf{F}) : \alpha^j = 0 \text{ for each } j = 0, ..., i\}$, where $j \in \mathbf{N_o}$. That is, $c_0(\mathbf{F}) \star K_i = K_i$, but $K_i \star c_0(\mathbf{F}) = K_{i-1}$ for each $i \in \mathbf{N_o}$, where $K_{-1} := c_0(\mathbf{F})$. The algebra $c_0(\mathbf{F})$ is the particular case of H, when $G_j = \{e\}$ for each $j \in \mathbf{N_o}$. We consider further H for non-trivial topological groups outlined above with $G_\infty := \bigcap_{j=0}^\infty G_j$ dense in each G_j.

22. Theorem. *If $\mathcal{F}$ is a maximal proper left or right ideal in H, then $H/\mathcal{F}$ is isomorphic as the nonassociative noncommutative algebra over $\mathbf{F}$ with $c_0(\mathbf{F})$.*

Proof. The ideal $\mathcal{F}$ is also the $\mathbf{F}$-linear subspace of H. In view of Theorem 7.12 [Roo78] a function $f : G_j \to \mathbf{F}$ is μ^j-integrable if and only if it satifies two properties: f is $Af(G,\mu^j)$-continuous and for each $\epsilon > 0$ the set $\{x : |f(x)|N_{\mu^j}(x) \ge \epsilon\}$ is $Af(G,\mu^j)$-compact and hence contained in $\{x : N_{\mu^j}(x) \ge \delta\}$ for some $\delta > 0$. Suppose, that there exists $j \in \mathbf{N_o}$ such that $f^j = 0$ for each $f \in \mathcal{F}$, then $f^i = 0$ for each $i \in \mathbf{N_o}$, since the space of bounded $\mathbf{F}$-valued continuous functions $C^0_b(G_\infty, \mathbf{F})$ on G_∞ is dense $H_j := \{f^j : f \in H\}$ and $C^0_b(G_\infty, \mathbf{F}) \cap F_j = \{0\}$ and $C^0_b(G_j, \mathbf{F})|_{G_{j+1}} \supset C^0_b(G_{j+1}, \mathbf{F})$. Therefore, $\mathcal{F}_j \ne \{0\}$ for each $j \in \mathbf{N_o}$, where $\mathcal{F}_j := \{f^j : f \in \mathcal{F}\}$, consequently, $\mathbf{F} \hookrightarrow F_j$ for each $j \in \mathbf{N_o}$. Since $\mathbf{F}$ is embeddable into each F_j, then there exists the embedding of $c_0(\mathbf{F})$ into $\mathcal{F}$, where $H_j := \{f^j : f \in H\}$, $\pi_j : H \to H_j$ are the natural projections.

The subalgebra $\mathcal{F}$ is closed in H, since H is the topological algebra and $\mathcal{F}$ is the maximal proper subalgebra. The space $H_\infty := \bigcap_{j \in \mathbf{N_o}} H_j$ is dense in each H_j.

If $\mathcal{F}_i = H_i$ for some $i \in \mathbf{N_o}$, then $\mathcal{F}_j = H_j$ for each $j \in \mathbf{N_o}$, since $C^0_b(G_\infty, \mathbf{F})$ is dense in each H_j and $C^0_b(G_j, \mathbf{F})|_{G_{j+1}} \supset C^0_b(G_{j+1}, \mathbf{F})$. The ideal $\mathcal{F}$ is proper, consequently, $\mathcal{F}_j \ne H_j$ as the $\mathbf{F}$-linear subspace for each $j \in \mathbf{N_o}$, where $\mathcal{F}_j = \pi_j(\mathcal{F})$.

There exist $\mathbf{F}$-linear continuous operators from $c_0(\mathbf{F})$ into $c_0(\mathbf{F})$ such that $x \mapsto (0, ..., 0, x^0, x^1, x^2, ...)$ with 0 as n coordinates at the beginning, $x \mapsto (x^n, x^{n+1}, x^{n+2}, ...)$ for $n \in \mathbf{N}$; $x \mapsto (x^{kl+\sigma_k(i)} : k \in \mathbf{N_o}, i \in (0, 1, ..., l-1))$, where $\mathbf{N} \ni l \ge 2$, $\sigma_k \in S_l$ are elements of the symmetric group S_l of the set $(0, 1, ..., l-1)$. Then $f \star (g \star h) + c_0(\mathbf{F})$ and $(f \star g) \star h + c_0(\mathbf{F})$ are considered as the same class, also $f \star g + c_0(\mathbf{F}) = g \star f + c_0(\mathbf{F})$ in $H/c_0(\mathbf{F})$, since $(f + c_0(\mathbf{F})) \star (g + c_0(\mathbf{F})) = f \star g + c_0(\mathbf{F})$ for each f, g and $h \in H$. Then $f \star (g \star h) + c_0(\mathbf{F})$ and $(f \star g) \star h + c_0(\mathbf{F})$ are considered as the same class for each $f, g, h \in \mathcal{F}$, also $f \star g + c_0(\mathbf{F}) = g \star f + c_0(\mathbf{F})$ in $\mathcal{F}/c_0(\mathbf{F})$, since $(f + c_0(\mathbf{F})) \star (g + c_0(\mathbf{F})) = f \star g + l_2(\mathbf{F}) \subset \mathcal{F}$ for each f and $g \in \mathcal{F}$. Therefore, the quotient algebras $H/c_0(\mathbf{F})$ and $\mathcal{F}/c_0(\mathbf{F})$ are the associative commutative Banach algebras.

From $\mu^i \in M_l(G_i, G_{i+1})$ it follows that for each open subset $W \ni e$, $W \subset G_i$ there exists a clopen subgroup $U \subset W$ such that $\mu^i(U) \ne 0$, since otherwise $\mu^i(zV) = 0$ for each $z \in G_{i+1}$ and each open $V \subset W$, hence $\mu^i(G_i) = 0$ contradicting supposition, that each μ^i is the probability measure.

Let us adjoin a unit to $H/c_0(\mathbf{F})$ and to $\mathcal{F}/c_0(\mathbf{F})$. There is satisfied the equality $Ch_{G_{i+1}} * Ch_{G_i} = Ch_{G_i}$. Let $U_{i,j}$ be a clopen subgroup in G_i, that is possible, since each G_i is ultrametrizable. Choose $U_{i,j}$ such that $U_{i,j+1} \subset U_{i,j} \cap G_{j+1}$ and $U_{i,j} \supset U_{i+1,j}$ for each i and j, $\bigcap_i U_{i,j} = e \in G_j$ for each j. Since $\mu^j(G_j) = 1$ and $\|G_j\|_{\mu^j} = 1$, then by induction $(i,j) \in \{(1,1),(1,2),...,(1,n),...;(2,1),(2,2),...,(2,n),...;...(m,1),(m,2),...,(m,n),...\}$, where $m,n \in \mathbf{N}$, there exists a family $\alpha_{i,j} \in \mathbf{F}$ and $\{U_{i,j} : i,j\}$ such that $\alpha_{i,j+1} Ch_{U_{i,j+1}} * \alpha_{i,j} Ch_{U_{i,j}} = \alpha_{i,j} Ch_{U_{i,j}}$ and $0 < |\alpha_{i,j}||\mu^i(U_{i,j})| \le 1$ for each i,j. Put $e_i := \{\alpha_{i,j} Ch_{U_{i,j}} : j \in \mathbf{N_o}\}$, then $e_i * e_i = e_i$ for each i. From the properties of $U_{i,j}$ it follows, that $span_{\mathbf{F}}\{e_i(z^{-1}g) : i \in \mathbf{N}, z \in G_\infty\}$ is dense in H, where $g := (g_j : g_j \in G_j \forall j \in \mathbf{N_o})$, $z^{-1}g = (z^{-1}g_j : j)$.

Consider the algebras $H/c_0(\mathbf{F}) =: A$ and $\mathcal{F}/c_0(\mathbf{F}) =: B$. The algebras A and B are commutative and associative. From the preceding proof it follows that $span_{\mathbf{F}}\{e_i(z^{-1}g) + c_0(\mathbf{F}) : i \in \mathbf{N},\ z \in G_\infty\}$ is dense in A, each $e_i(z^{-1}g) + c_0(\mathbf{F})$ is the idempotent element in A. Therefore, by Theorem 6.12 [Roo78] A is the C-algebra. By the definition this means, that there exists a locally compact zero-dimensional Hausdorff space X such that A is isomorphic with $C_\infty(X,\mathbf{F})$, where $C_\infty(X,\mathbf{F})$ is the subspace of all $f \in C_b(X,\mathbf{F})$ for which for each $\epsilon > 0$ there exists a compact subset $X_{\epsilon,f}$ of X with $|f(x)| < \epsilon$ for each $x \in X \setminus X_{\epsilon,f}$. In accordance with Theorem 6.3 [Roo78] each maximal ideal $\mathcal{B}$ of $C_\infty(X,\mathbf{F})$ has the form $\mathcal{B} = \{f \in C_\infty(X,\mathbf{F}) : f(z_0) = 0\}$, where z_0 is a marked point in X. On the other hand, as it was proved above $\mathcal{F}_j \ne H_j$ for each $j \in \mathbf{N_o}$, hence there exists the following embedding $c_0(\mathbf{F}) \hookrightarrow (H/\mathcal{F})$ and $(H/\mathcal{F})/c_0(\mathbf{F})$ is isomorphic with $(H/c_0(\mathbf{F}))/(\mathcal{F}/c_0(\mathbf{F}))$. Therefore, $H/\mathcal{F}$ is isomorphic with $c_0(\mathbf{F})$.

23. Comments. Another methods of construction of isometrical representations of topological totally disconnected groups which may be nonlocally compact with the help of quasi-invariant $\mathbf{F}$-valued measures were given in [Lud98b, Lud02b, Lud00a, Lud99t, Lud01s, Lud0348, Lud01f, LD03] and references therein.

Using results of this article it is possible to make further investigations of nonlocally compact totally disconnected topological groups, their structures and representations, measurable operators in Banach spaces over non-Archimedean fields, apply this for the development of non-Archimedean quantum mechanics and quantum field theory, quantum gravity, superstring theory and gauge theory, etc.

References

[Ami64] Amice, Y.: Interpolation p-adique. Bull. Soc. Math. France, **92**, 117–180 (1964)

[ADV88] Aref'eva, I.Ya., Dragovich, B., Volovich, I.V.: On the p-adic summability of the anharmonic oscillator. Phys. Lett., **B 200**, 512–514 (1988)

[BGR84] Bosch, S., Guntzer, U., Remmert, R.: Non-Archimedean analysis. Springer, Berlin (1984)

[Bou76] Bourbaki, N.: Lie groups and algebras. Mir, Moscow (1976)

[Bou63-69] Bourbaki, N.: Intégration. Livre VI. Fasc. XIII, XXI, XXIX, XXXV. Ch. 1–9. Hermann, Paris (1965, 1967, 1963, 1969).

[Cas02] Castro, C.: Fractal strings as an alternative justification for El Naschie's cantorian spacetime and the fine structure constants. Chaos, Solitons and Fractals, **14**, 1341–1351 (2002)

[Dia84] Diarra, B.: Ultraproduits ultrametriques de corps values. Ann. Sci. Univ. Clermont II, Sér. Math., **22**, 1-37 (1984)

[DD00] Djordjević, G.S., Dragovich, B.: p-Adic and adelic harmonic oscillator with a time-dependent frequency. Theor. and Math. Phys., **124: 2**, 1059-1067 (2000)

[Eng86] Engelking, R.: General topology. Mir, Moscow (1986)

[Esc95] Escassut, A.: Analytic elements in p-adic analysis. World Scientific, Singapore (1995)

[FD88] Fell, J.M.G., Doran, R.S.: Representations of $*$-algebras, locally compact groups, and Banach $*$-algebraic bundles. Acad. Press, Boston (1988)

[FP81] Fresnel, J., Put, M. van der.: Géométrie analytique rigide et applications. Birkhäuser, Boston (1981)

[Gru66] Gruson, L.: Théorie de Fredholm p-adique. Bull. Soc. Math. France, **94**, 67-95 (1966)

[HR79] Hewitt, E., Ross, K.A.: Abstract harmonic analysis. Second ed. Springer, Berlin (1979)

[Jan98] Jang, Y.: Non-Archimedean quantum mechanics. Tohoku Math. Publ. N **10** (1998)

[Khr90] Khrennikov, A.Yu.: Mathematical methods of non-Archimedean physics. Russ. Math. Surv., **45: 4**, 79–110 (1990)

[Kob77] Koblitz, N.: p-adic numbers, p-adic analysis and zeta functions. Springer-Verlag, New York, 1977.

[Lud96] Ludkovsky, S.V.: Measures on groups of diffeomorphisms of non-Archimedean Banach manifolds. Russ. Math. Surv., **51: 2**, 338–340 (1996)

[Lud96c] Ludkovsky, S.V.: Quasi-invariant and pseudo-differentiable measures on a non-Archimedean Banach space. I, II. Los Alamos Preprints **math.GM/0106169** and **math.GM/0106170** (http://xxx.lanl.gov/; earlier version: ICTP **IC/96/210**, October 1996, 50 p.p.; http://www.ictp.trieste.it/), II to appear in J. Math. Sci.

[Lud98b] Ludkovsky, S.V.: Irreducible unitary representations of non-Archimedean groups of diffeomorphisms. Southeast Asian Bull. of Math., **22**, 419–436 (1998)

[Lud98s] Ludkovsky, S.V.: Quasi-invariant measures on non-Archimedean semigroups of loops. Russ. Math. Surv., **53: 3**, 633–634 (1998)

[Lud99a] Ludkovsky, S.V.: Properties of quasi-invariant measures on topological groups and associated algebras. Annales Math. B. Pascal, **6: 1**, 33–45 (1999)

[Lud99t] Ludkovsky, S.V.: Measures on groups of diffeomorphisms of non-Archimedean manifolds, representations of groups and their applications. Theoret. and Math. Phys., **119: 3**, 698–711 (1999)

[Lud00a] Ludkovsky, S.V.: Quasi-invariant measures on non-Archimedean groups and semigroups of loops and paths, their representations. I, II. Annales Math. B. Pascal, **7: 2**, 19–53, 55–80 (2000)

[Lud00f] Ludkovsky, S.V.: Non-Archimedean polyhedral decompositions of ultrauniform spaces. Fundam. i Prikl. Math. **6: 2**, 455–475 (2000)

[Lud01f] Ludkovsky, S.V.: Stochastic processes on groups of diffeomorphisms and loops of real, complex and non-Archimedean manifolds. Fundam. i Prikl. Math., **7: 4**, 1091-1105 (2001)

[Lud01s] Ludkovsky, S.V.: Representations of topological groups generated by Poisson measures. Russ. Math. Surv., **56: 1**, 169–170 (2001)

[Lud02a] Ludkovsky, S.V.: Quasi-invariant and pseudo-differentiable real-valued measures on a non-Archimedean Banach space. Analysis Math., **28**, 287–316 (2002)

[Lud02b] Ludkovsky, S.V.: Poisson measures for topological groups and their representations. Southeast Asian Bull. Math., **25: 4**, 653–680 (2002)

[Lud0321] Ludkovsky, S.V.: Stochastic processes on non-Archimedean Banach spaces. Int. J. of Math. and Math. Sci., **2003: 21**, 1341–1363 (2003)

[Lud0341] Ludkovsky, S.V.: Stochastic antiderivational equations on non-Archimedean Banach spaces. Int. J. of Math. and Math. Sci., **2003: 41**, 2587–2602 (2003)

[Lud0348] Ludkovsky, S.V.: Stochastic processes on totally disconnected topological groups. Int. J. of Math. and Math. Sci., **2003: 48**, 3067–3089 (2003)

[Lud02j] Ludkovsky, S.V.: Quasi-invariant and pseudo-differentiable measures on non-Archimedean Banach spaces with values in non-Archimedean fields. J. Math. Sci., **112**, 54 pages (2002)

[Lud03s2] Ludkovsky, S.V.: Quasi-invariant and pseudo-differentiable measures on non-Archimedean Banach spaces. Russ. Math. Surv., **58: 2**, 167–168 (2003)

[Lud03s6] Ludkovsky, S.V.: A structure of groups of diffeomorphisms of non-Archimedean manifolds. Russ. Math. Surv., **58: 6**, 155–156 (2003)

[Lud03b] Ludkovsky, S.V.: A structure and representations of diffeomorphism groups of non-Archimedean manifolds. Southeast Asian Bull. of Math., **26**, 975-1004 (2003)

[LD03] Ludkovsky, S., Diarra, B.: Profinite and finite groups associated with loop and diffeomorphism groups of non-Archimedean manifolds. Int. J. Math. and Math. Sci., **2003: 42**, 2673–2688 (2003)

[LK02] Ludkovsky, S.V., Khrennikov, A.: Stochastic processes on non-Archimedean spaces with values in non-Archimedean fields. Markov Processes and Related Fields, **8**, 1-34 (2002)

[MS63] Monna, A.P., Springer, T.A.: Integration non-archimedienne. Indag. Math., **25**, 634-653 (1963)

[Nai68] Naimark, M.A.: Normed rings. Nauka, Moscow (1968)

[NB85] Narici, L., Beckenstein, E.: Topological vector spaces. Marcel Dekker Inc., New York (1985)

[Put68] Put, M. van der: The ring of bounded operators on a non-Archimedean normed linear space. Indag. Math. **71: 3**, 260–264 (1968)

[Roo78] Rooij, A.C.M. van.: Non-Archimedean functional analysis. Marcel Dekker Inc., New York (1978)

[Sch84] Schikhof, W.H.: Ultrametric calculus. Camb. Univ. Press, Cambridge (1984)

[Sch89] Schikhov, W.H.: On p-adic compact operators. Report **8911**. Dept. Math. Cath. Univ., Nijmegen, The Netherlands (1989)

[Sch71] Schikhof, W.H.: A Radon-Nikodym theorem for non-Archimedean integrals and absolutely continuous measures on groups. Indag. Math. Ser. A, **33: 1**, 78–85 (1971)

[VV89] Vladimirov, V.S., Volovich, I.V. Comm. Math. Phys., **123** , 659–676 (1989)

[VVZ94] Vladimirov, V.S., Volovich, I.V., Zelenov, E.I.: p-Adic analysis and mathematical physics. Fiz.-Mat. Lit., Moscow (1994)

[Wei73] Weil, A.: Basic number theory. Springer, Berlin (1973)